

Bayesian state-space models for affective dynamics

Tom Lodewyckx, Francis Tuerlinckx, Peter Kuppens
Quantitative psychology & individual differences
KULeuven

Nicholas Allen
University of Melbourne

Lisa Sheeber
Oregon Research Institute

OUTLINE


- 1 State-space models
- 2 Bayesian estimation
- 3 Application
- 4 Discussion

OUTLINE

- 1 **State-space models**
 - *Linear dynamical systems*
- 2 Bayesian estimation
- 3 Application
- 4 Discussion

Linear dynamical system theory

- *Inputs (observed)*: control / noise / demographic variables
- *System (unobserved)*: latent states with dynamical dependencies
- *Outputs (observed)*: measurements of interest



OUTLINE

1 State-space models

- Linear dynamical systems
- *Multivariate linear Gaussian state-space model*

2 Bayesian estimation

3 Application

4 Discussion

Application fields

- Engineering
- Econometrics
- Ecological research

Application fields

- Engineering
- Econometrics
- Ecological research

Model structure

- Transition equation: latent dynamical system
- Observation equation: observed variables

Transition equation

$$Y_t \sim \text{Gaussian}(\Psi\Theta_t + \Gamma X_t, \Sigma_\epsilon)$$

$$\Theta_t \sim \text{Gaussian}(\Phi\Theta_{t-1} + \Delta Z_t, \Sigma_\eta)$$


- Θ_t = states
- Θ_{t-1} = states at previous observation moment
- Z_t = state covariates

Transition equation

$$Y_t \sim \text{Gaussian}(\Psi\Theta_t + \Gamma X_t, \Sigma_\epsilon)$$

$$\Theta_t \sim \text{Gaussian}(\Phi\Theta_{t-1} + \Delta Z_t, \Sigma_\eta)$$

- Θ_t = states
- Θ_{t-1} = states at previous observation moment
- Z_t = state covariates
- Φ = transition matrix (autocorrelations & cross-lagged relations)
- Δ = state covariate regression coefficient matrix
- Σ_η = innovation variance/covariance matrix



Observation equation

$$Y_t \sim \text{Gaussian}(\Psi\Theta_t + \Gamma X_t, \Sigma_\epsilon)$$

$$\Theta_t \sim \text{Gaussian}(\Phi\Theta_{t-1} + \Delta Z_t, \Sigma_\eta)$$


- Y_t = observations
- Θ_t = states
- X_t = observation covariates


Observation equation

$$Y_t \sim \text{Gaussian}(\Psi\Theta_t + \Gamma X_t, \Sigma_\epsilon)$$

$$\Theta_t \sim \text{Gaussian}(\Phi\Theta_{t-1} + \Delta Z_t, \Sigma_\eta)$$

- Y_t = observations
- Θ_t = states
- X_t = observation covariates
- Ψ = design matrix (mapping $\Theta_t \rightarrow Y_t$)
- Γ = observation covariate regression coefficient matrix
- Σ_ϵ = observation error variance/covariance matrix





OUTLINE

1 State-space models

- Linear dynamical systems
- Multivariate linear Gaussian state-space model
- *Regime-switching MLGSS model*

2 Bayesian estimation

3 Application

4 Discussion

Regime-switching MLGSS model

$$Y_t \sim \text{Gaussian}(\Psi_t \Theta_t + \Gamma_t X_t, \Sigma_{\epsilon t})$$

$$\Theta_t \sim \text{Gaussian}(\Phi_t \Theta_{t-1} + \Delta_t Z_t, \Sigma_{\eta t})$$

Regime-switching MLGSS model

$$Y_t \sim \text{Gaussian}(\Psi_t \Theta_t + \Gamma_t X_t, \Sigma_{\epsilon t})$$

$$\Theta_t \sim \text{Gaussian}(\Phi_t \Theta_{t-1} + \Delta_t Z_t, \Sigma_{\eta t})$$

$$\begin{pmatrix} \Psi_t, \Gamma_t \\ \Phi_t, \Delta_t \end{pmatrix} = \begin{cases} \begin{pmatrix} \Psi_1, \Gamma_1 \\ \Phi_1, \Delta_1 \end{pmatrix} & \text{if } R_t = 1 \\ \dots \\ \begin{pmatrix} \Psi_r, \Gamma_r \\ \Phi_r, \Delta_r \end{pmatrix} & \text{if } R_t = r \end{cases}$$

OUTLINE

- 1 State-space models
- 2 **Bayesian estimation**
- 3 Application
- 4 Discussion

Gibbs sampling

- Bayesian estimation method, iterative Monte Carlo simulation
- Each parameter has conditional sampling distribution
- Blocking of parameters efficient for certain models
(no WinBUGS ☹)

Gibbs sampling

- Bayesian estimation method, iterative Monte Carlo simulation
- Each parameter has conditional sampling distribution
- Blocking of parameters efficient for certain models
(no WinBUGS ☹)


Blocked Gibbs sampler

- 1 Sampling distribution Θ
- 2 Sampling distribution $\Psi_1, \dots, \Psi_r, \Gamma_1, \dots, \Gamma_r, \Sigma_\epsilon$
- 3 Sampling distribution $\Phi_1, \dots, \Phi_r, \Delta_1, \dots, \Delta_r, \Sigma_\eta$


Blocked Gibbs sampler

- 1 Sampling distribution Θ
→ Forward-filtering backward sampling
(Carter & Kohn, 1994, 1996)
- 2 Sampling distribution $\Psi_1, \dots, \Psi_r, \Gamma_1, \dots, \Gamma_r, \Sigma_\epsilon$
- 3 Sampling distribution $\Phi_1, \dots, \Phi_r, \Delta_1, \dots, \Delta_r, \Sigma_\eta$

\Rightarrow Kalman filter \Rightarrow



\Rightarrow Kalman filter \Rightarrow



\Leftarrow Backward-sampling \Leftarrow

Blocked Gibbs sampler

- 1 Sampling distribution Θ
- 2 Sampling distribution $\Psi_1, \dots, \Psi_r, \Gamma_1, \dots, \Gamma_r, \Sigma_\epsilon$
→ Method of decomposition
(Zellner, 1962)
- 3 Sampling distribution $\Phi_1, \dots, \Phi_r, \Delta_1, \dots, \Delta_r, \Sigma_\eta$
→ Method of decomposition
(Zellner, 1962)

Parameters observation equation

$\Sigma_{\epsilon} \rightarrow$ Inverse Wishart

Parameters observation equation

$\Sigma_{\epsilon} \rightarrow$ Inverse Wishart

$\Psi_1, \dots, \Psi_r \mid \Sigma_{\epsilon} \rightarrow$ Multivariate Gaussian

Parameters observation equation

$$\Sigma_{\epsilon} \rightarrow \text{Inverse Wishart}$$

$$\Psi_1, \dots, \Psi_r \mid \Sigma_{\epsilon} \rightarrow \text{Multivariate Gaussian}$$

$$\Delta_1, \dots, \Delta_r \mid \Sigma_{\epsilon}, \Psi_1, \dots, \Psi_r \rightarrow \text{Multivariate Gaussian}$$

Parameters observation equation

$$\Sigma_{\epsilon} \rightarrow \text{Inverse Wishart}$$

$$\Psi_1, \dots, \Psi_r \mid \Sigma_{\epsilon} \rightarrow \text{Multivariate Gaussian}$$

$$\Delta_1, \dots, \Delta_r \mid \Sigma_{\epsilon}, \Psi_1, \dots, \Psi_r \rightarrow \text{Multivariate Gaussian}$$

Parameters transition equation

$$\Sigma_{\eta} \rightarrow \text{Inverse Wishart}$$

$$\Phi_1, \dots, \Phi_r \mid \Sigma_{\eta} \rightarrow \text{Multivariate Gaussian}$$

$$\Gamma_1, \dots, \Gamma_r \mid \Sigma_{\eta}, \Phi_1, \dots, \Phi_r \rightarrow \text{Multivariate Gaussian}$$

OUTLINE

- 1 State-space models
- 2 Bayesian estimation
- 3 **Application**
- 4 Discussion

Emotion regulation

- High: emotions are adapted to the social context

Low: emotions are insensitive to the context

Emotion regulation

- High: emotions are adapted to the social context
Low: emotions are insensitive to the context
- Baseline respiratory sinus arrhythmia (QRSA)
Butler, Wilhelm & Gross (2006); Gyurak & Ayduk (2008)

Emotion regulation

- High: emotions are adapted to the social context
Low: emotions are insensitive to the context
- Baseline respiratory sinus arrhythmia (QRSA)
Butler, Wilhelm & Gross (2006); Gyurak & Ayduk (2008)

Emotional inertia

- High: emotions don't change much over time
Low: emotions constantly fluctuate over time

Emotion regulation

- High: emotions are adapted to the social context
Low: emotions are insensitive to the context
- Baseline respiratory sinus arrhythmia (QRSA)
Butler, Wilhelm & Gross (2006); Gyurak & Ayduk (2008)

Emotional inertia

- High: emotions don't change much over time
Low: emotions constantly fluctuate over time
- Autocorrelation of emotional process (ϕ_{em})
Kuppens, Allen & Sheeber (submitted)

Oregon adolescent interaction data

- 69 normal & 72 depressed adolescents
- Adolescent, father and mother in laboratory
- Problem solving task (e.g. pocket money)
- One measure/second during 18 minutes ($n = 1080$)

Oregon adolescent interaction data

- 69 normal & 72 depressed adolescents
- Adolescent, father and mother in laboratory
- Problem solving task (e.g. pocket money)
- One measure/second during 18 minutes ($n = 1080$)

Measurements

- QRSA (emotion regulation)

Oregon adolescent interaction data

- 69 normal & 72 depressed adolescents
- Adolescent, father and mother in laboratory
- Problem solving task (e.g. pocket money)
- One measure/second during 18 minutes ($n = 1080$)

Measurements

- QRSA (emotion regulation)
- HR_t (heart rate, “Fight-or-flight”)
- RSA_t (respiratory sinus arrhythmia, “Rest-and-digest”)

Oregon adolescent interaction data

- 69 normal & 72 depressed adolescents
- Adolescent, father and mother in laboratory
- Problem solving task (e.g. pocket money)
- One measure/second during 18 minutes ($n = 1080$)

Measurements

- QRSA (emotion regulation)
- HR_t (heart rate, “Fight-or-flight”)
- RSA_t (respiratory sinus arrhythmia, “Rest-and-digest”)
- $AdAnger_t$ (anger of adolescent)

Basic MLGSS Model

- $[Y_{\text{HR}} \ Y_{\text{RSA}}]_t \sim \text{Gaussian} (I_2[\Theta_{\text{HR}} \ \Theta_{\text{RSA}}]_t, \Sigma_{\epsilon})$

Basic MLGSS Model

- $[Y_{\text{HR}} \ Y_{\text{RSA}}]_t \sim \text{Gaussian} (I_2[\Theta_{\text{HR}} \ \Theta_{\text{RSA}}]_t, \Sigma_{\epsilon})$
 $[\Theta_{\text{HR}} \ \Theta_{\text{RSA}}]_t \sim \text{Gaussian} (\Phi[\Theta_{\text{HR}} \ \Theta_{\text{RSA}}]_{t-1}, \Sigma_{\eta})$

Basic MLGSS Model

- $[Y_{\text{HR}} \ Y_{\text{RSA}}]_t \sim \text{Gaussian} (I_2[\Theta_{\text{HR}} \ \Theta_{\text{RSA}}]_t, \Sigma_{\epsilon})$
- $[\Theta_{\text{HR}} \ \Theta_{\text{RSA}}]_t \sim \text{Gaussian} (\Phi[\Theta_{\text{HR}} \ \Theta_{\text{RSA}}]_{t-1}, \Sigma_{\eta})$

Results

	Normal	Depressed
$\hat{\phi}_{\text{HR}}$.79 (.08)	.81 (.08)

Basic MLGSS Model

- $[Y_{\text{HR}} \ Y_{\text{RSA}}]_t \sim \text{Gaussian} (I_2[\Theta_{\text{HR}} \ \Theta_{\text{RSA}}]_t, \Sigma_{\epsilon})$
- $[\Theta_{\text{HR}} \ \Theta_{\text{RSA}}]_t \sim \text{Gaussian} (\Phi[\Theta_{\text{HR}} \ \Theta_{\text{RSA}}]_{t-1}, \Sigma_{\eta})$

Results

	Normal	Depressed
$\hat{\phi}_{\text{HR}}$.79 (.08)	.81 (.08)
$r(\hat{\phi}_{\text{HR}}, \text{QRSA})$	-.28	-.64

Regime-switching MLGSS Model

- $[Y_{\text{HR}} \ Y_{\text{RSA}}]_t \sim \text{Gaussian}(\mathbf{I}_2[\Theta_{\text{HR}} \ \Theta_{\text{RSA}}]_t, \Sigma_{\epsilon})$
- $[\Theta_{\text{HR}} \ \Theta_{\text{RSA}}]_t \sim \text{Gaussian}(\Phi_t[\Theta_{\text{HR}} \ \Theta_{\text{RSA}}]_{t-1}, \Sigma_{\eta})$
- $$\Phi_t = \begin{cases} \Phi_{[\text{no anger}]} & \text{if } \text{AdAnger}_t = 0 \\ \Phi_{[\text{anger}]} & \text{if } \text{AdAnger}_t = 1 \end{cases}$$

Regime-switching MLGSS Model

- $[Y_{\text{HR}} \ Y_{\text{RSA}}]_t \sim \text{Gaussian}(\mathbf{I}_2[\Theta_{\text{HR}} \ \Theta_{\text{RSA}}]_t, \Sigma_{\epsilon})$
- $[\Theta_{\text{HR}} \ \Theta_{\text{RSA}}]_t \sim \text{Gaussian}(\Phi_t[\Theta_{\text{HR}} \ \Theta_{\text{RSA}}]_{t-1}, \Sigma_{\eta})$
- $\Phi_t = \begin{cases} \Phi_{[\text{no anger}]} & \text{if } \text{AdAnger}_t = 0 \\ \Phi_{[\text{anger}]} & \text{if } \text{AdAnger}_t = 1 \end{cases}$

Results

	Normal	Depressed
$\hat{\phi}_{\text{HR}[\text{no anger}]}$.79 (.08)	.80 (.08)
$\hat{\phi}_{\text{HR}[\text{anger}]}$.73 (.16)	.78 (.11)

Regime-switching MLGSS Model

- $[Y_{\text{HR}} \ Y_{\text{RSA}}]_t \sim \text{Gaussian} (I_2[\Theta_{\text{HR}} \ \Theta_{\text{RSA}}]_t, \Sigma_{\epsilon})$
- $[\Theta_{\text{HR}} \ \Theta_{\text{RSA}}]_t \sim \text{Gaussian} (\Phi_t[\Theta_{\text{HR}} \ \Theta_{\text{RSA}}]_{t-1}, \Sigma_{\eta})$
- $\Phi_t = \begin{cases} \Phi_{[\text{no anger}]} & \text{if } \text{AdAnger}_t = 0 \\ \Phi_{[\text{anger}]} & \text{if } \text{AdAnger}_t = 1 \end{cases}$

Results

	Normal	Depressed
$\hat{\phi}_{\text{HR}[\text{no anger}]}$.79 (.08)	.80 (.08)
$\hat{\phi}_{\text{HR}[\text{anger}]}$.73 (.16)	.78 (.11)
$r(\hat{\phi}_{\text{HR}[\text{no anger}]}, \text{QRSA})$	-.26	-.67
$r(\hat{\phi}_{\text{HR}[\text{anger}]}, \text{QRSA})$	-.10	-.65

Summary

- No difference in heart rate inertia between normal and depressed subjects

Summary

- No difference in heart rate inertia between normal and depressed subjects
- Negative relation between heart rate inertia and QRSA:
The higher (lower) the degree of emotion regulation,
the weaker (stronger) the emotional inertia

Summary

- No difference in heart rate inertia between normal and depressed subjects
- Negative relation between heart rate inertia and QRSA:
The higher (lower) the degree of emotion regulation,
the weaker (stronger) the emotional inertia
(asymmetry normal & depressed?)

OUTLINE

- 1 State-space models
- 2 Bayesian estimation
- 3 Application
- 4 **Discussion**

Modeling issues

- Hierarchical modeling (individual & group differences)
- Prior distributions
- Latent regime variable
- Hypothesis testing

Modeling issues

- Hierarchical modeling (individual & group differences)
- Prior distributions
- Latent regime variable
- Hypothesis testing

Conceptual issues

- Factor structures in design matrix
- Non-Gaussian errors
- Lagging factor

Conclusion

- 1 Framework useful for latent psychophysiological dynamics

Conclusion

- 1 Framework useful for latent psychophysiological dynamics
- 2 Bayesian estimation using MCMC methods

Conclusion

- 1 Framework useful for latent psychophysiological dynamics
- 2 Bayesian estimation using MCMC methods
- 3 Hierarchical model more elegant

Conclusion

- 1 Framework useful for latent psychophysiological dynamics
- 2 Bayesian estimation using MCMC methods
- 3 Hierarchical model more elegant
- 4 Various other model extensions to be considered

Conclusion

- 1 Framework useful for latent psychophysiological dynamics
- 2 Bayesian estimation using MCMC methods
- 3 Hierarchical model more elegant
- 4 Various other model extensions to be considered

Questions?